



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the vertical distance of the dove from A . This may be represented by $c\theta$. It follows that $BP = \sqrt{a^2\theta^2 + c^2\theta^2} = (a^2 + c^2)^{\frac{1}{2}}\theta$.

The co-ordinates of P are :

$$x = a \cos \theta + (a^2 + c^2)^{\frac{1}{2}} \theta \sin \theta,$$

$$y = a \sin \theta - (a^2 + c^2)^{\frac{1}{2}} \theta \cos \theta,$$

$$z = c\theta, \text{ the axis of } z \text{ coinciding with the cylinder's axis. } \therefore ds^2 = dx^2 + dy^2 + dz^2 = \left[2a^2 + 2c^2 - 2a(a^2 + c^2)^{\frac{1}{2}} + (a^2 + c^2)\theta^2 \right] d\theta^2.$$

$$\therefore s = \int \left[2a^2 + 2c^2 - 2a(a^2 + c^2)^{\frac{1}{2}} + (a^2 + c^2)\theta^2 \right]^{\frac{1}{2}} d\theta = \sqrt{a^2 + c^2} \left\{ \frac{\theta}{2} \left(\frac{2a^2 + 2c^2 - 2a(a^2 + c^2)^{\frac{1}{2}}}{a^2 + c^2} + \theta^2 \right)^{\frac{1}{2}} + \frac{a^2 + c^2 - a\sqrt{a^2 + c^2}}{a^2 + c^2} \log \left[\theta + \left(\frac{2a^2 + 2c^2 - 2a\sqrt{a^2 + c^2}}{a^2 + c^2} + \theta^2 \right)^{\frac{1}{2}} \right] \right\}.$$

$$a=1, c=\frac{1}{2\pi}, \text{ and the } \theta\text{-limits are } 40\pi \text{ and } 0.$$

$$\therefore s = 7995 \text{ feet, approximately.}$$

[*C. E. Myers* solves for case I. and gets 7895.517768 feet. *Prof. Zerr* solves by a different method and gets 8023.754 feet and 8124.910 + feet, for his results.]

PROBLEMS.

12. Proposed by **ISAAC L. BEVERAGE**, Monterey, Virginia.

Given the equations $2z^3 = x + 3z$ and $5z^2 = y + 2z$. To find $\frac{dy}{dx}$ for $x=0$.

13. Proposed by **J. A. CALDERHEAD**, B. Sc., Superintendent of Schools, Lima, Ohio.

A steamer whose course is due west and speed 10 knots is sighted by another steamer going at 8 knots; what course must the latter steer, so as to cross the track of the former at the least possible distance from her?

14. Proposed by **J. F. W. SCHEFFER**, A. M., Hagerstown, Maryland.

Right triangles are inscribed in a circle whose centre $= (a, b)$, and radius $= c$. If one of the legs passes through a fixed point, prove that $c^2(x^2 + y^2) = (a^2 + b^2 - c^2 - ax - by)^2$ is the curve to which the other leg is always tangent; the fixed point being the origin of the co-ordinates.

15. Proposed by **CHARLES E. MYERS**, Canton, Ohio.

From a given quantity of material a cylindrical cup with circular bottom and open top is to be made, the cup to contain the greatest amount. What must be its dimensions?

Solutions to these problems should be received on or before May 1st.